

An Estimate of the Branching Fraction of $\tau \rightarrow \pi\eta'\nu_\tau$

S. Nussinov^{1,2} and A. Soffer¹

¹*Tel Aviv University, Tel Aviv, 69978, Israel*

²*Schmid College of Science, Chapman University, Orange, CA 92866, USA*

(Dated: July 21, 2009)

We calculate the expected branching fraction of the second-class-current decay $\tau \rightarrow \pi\eta'\nu_\tau$, motivated by a recent experimental upper-limit determination of this quantity. The largest contribution to the branching fraction is due to the intermediate $a_0(980)$ scalar meson, assuming it is a $\bar{u}d$ state. Smaller contributions arise from $a_0(1450)$, $\rho(770)$, and $\rho(1450)$. Our calculated values are substantially below the experimental upper limit, and are smaller still if the $a_0(980)$ is a four-quark state, as often suggested. Thus, a precise measurement or tight upper limit has the potential to determine the nature of the $a_0(980)$, as well as search for new scalar interactions.

PACS numbers:

I. INTRODUCTION

In a recent paper [1], we considered the branching fraction of the isospin-violating decay $\tau \rightarrow \pi\eta\nu_\tau$. We found an expected branching fraction of

$$\mathcal{B} \equiv \mathcal{B}(\tau \rightarrow \pi\eta\nu_\tau) = (0.3 - 1.0) \times 10^{-5}, \quad (1)$$

in rough agreement with a detailed chiral-perturbation-theory calculation [2] and other evaluations [3], which yielded central values in the range

$$\mathcal{B} = (1.2 - 1.6) \times 10^{-5}. \quad (2)$$

The experimental bound on this branching fraction, $\mathcal{B} < 1.4 \times 10^{-4}$ [4], was obtained by CLEO with an e^+e^- -collision data sample of 3.5 fb^{-1} , a fraction of a percent of currently available integrated luminosity. The only related high-luminosity measurement is a stringent BABAR upper limit on the branching fraction of $\tau \rightarrow \pi\eta'\nu_\tau$ [5],

$$\mathcal{B}' \equiv \mathcal{B}(\tau \rightarrow \pi\eta'\nu_\tau) < 7.2 \times 10^{-6} \quad @90\% \text{ CL}, \quad (3)$$

obtained with an integrated luminosity of 384 fb^{-1} .

The fact that the experimental limit is lower than the results summarized in Eq. (2) raises the question of a possible discrepancy between theory and experiment. Therefore, our goal in this article is to calculate the expected value of $\mathcal{B}'$ and compare it to the experimental limit. We adapt the methods used in Ref. [1] to the present case, noting that a chiral-perturbation-theory calculation of this process, as performed for $\tau \rightarrow \pi\eta\nu_\tau$ by Neufeld and Rupertsberger [2], would be very useful.

First, we note several similarities and differences between the calculations of $\mathcal{B}'$ and $\mathcal{B}$:

- The $\bar{u}u + \bar{d}d$ fraction of the wave function which, unlike the $\bar{s}s$ and gg parts, contributes to the decay amplitude, may be smaller for the η' . While it appears that the magnitude of the $\bar{s}s$ part in relation to that of the light quarks is very similar for both states, the current estimate of the gg fraction of the wave function, Z_{gg} , is $|Z_{gg}|^2 = 0.3 \pm 0.2$ [6].

In our calculations we take $Z_{gg} = 0$, as this yields the most conservative limits on $\mathcal{B}'$, and since the modification for finite values of Z_{gg} is straightforward.

- Calculations of $\mathcal{B}$ in Refs. [1, 2, 3] rely on extrapolations utilizing intermediate, low-mass $J^{PC} = 1^{--}$ and 0^{++} hadrons. Obvious intermediate states for the decay $\tau \rightarrow \pi\eta\nu_\tau$ are the ground-state mesons $\rho(770)$ and $a_0(980)$. In the case of $\tau \rightarrow \pi\eta'\nu_\tau$, these are off-shell processes, and the contributions of these resonances are suppressed. On the other hand, we do have now on-shell decays involving the next 1^{--} and 0^{++} states. These are the $\rho' \equiv \rho(1450)$ and $a_0(1450)$, which contribute to the P - and S -wave components of the decay, respectively.
- The ρ and ρ' vectors are the quark-model $\bar{u}d$, S -wave 1^{--} ground state and first radial excitation, respectively. However, the theoretical assignment of the $a_0(980)$ (and, consequently, that of the $a_0(1450)$ as well) is ambiguous, generating the largest uncertainty in both $\mathcal{B}$ and $\mathcal{B}'$. Conversely, information on these branching fractions can help resolve the longstanding dilemma of the “ $\bar{K}K$ -threshold” state $a_0(980)$. The significant branching fractions of $a_0(980)$ and $f_0(980)$ decays to $\bar{K}K$, despite the very small phase space, seem inconsistent with these mesons being the ground states of the quark-model scalar nonet, motivating a four-quark ($\bar{u}d\bar{s}s$) interpretation [7]. In this case, the $\bar{u}d$ scalar ground state should most likely be identified with $a_0(1450)$. However, this would make the scalar 190 MeV heavier than the axial vector state $a_1(1260)$, implying a pattern of $L \cdot S$ splitting different from what is observed in any other $L = 1$, $\bar{q}q'$ system. The more appealing possibility, namely, that the two 980-MeV states are indeed just $\bar{u}d$ states, may have been partially resurrected in recent work [8], in which 'tHooft's $\bar{u}u\bar{d}\bar{d}\bar{s}s$ six-quark vertex was utilized to admix the 2- and 4- quark states.

The plan of this note is as follows. As we did in Ref. [1], we discuss separately our estimates of the P - and S -wave contributions to $\mathcal{B}'$. In Sec II we present the more robust results for the P -wave part, calculating upper bounds on the contributions of the ρ and ρ' using recently published experimental data involving η' and τ decays. In Sec III we present the less clear-cut estimate of the S -wave component. This contribution depends most strongly on whether the $a_0(980)$ is a 4-quark state or the $\bar{u}d$ ground state. In any event, our predictions for $\mathcal{B}(\tau \rightarrow \pi\eta'\nu_\tau)$ lie significantly below the *BABAR* limit [5]. A brief summary and future outlook are given in Sec IV.

II. THE $L = 1$ CONTRIBUTION

In Ref. [1], we obtained the $L = 1$ contribution to $\mathcal{B}$ assuming that it was dominated by the ρ , an assumption justified by the large branching fraction $\mathcal{B}(\tau \rightarrow \rho\nu_\tau)$. We compared this branching fraction to $\mathcal{B}$ using the ratio of coupling constants $g_{\eta\rho\pi}/g_{\rho\pi\pi}$, where $g_{\rho\pi\pi}$ was related to the width of the ρ , and $g_{\eta\rho\pi}$ was obtained by analyzing the Dalitz-plot distribution of the decay $\eta \rightarrow \pi^+\pi^-\pi^0$, taking the scalar contribution to $\eta \rightarrow \pi^+\pi^-\pi^0$ from $\mathcal{B}(\eta \rightarrow \pi^0\pi^0\pi^0)$.

This procedure is not directly applicable to $\mathcal{B}'$, since there is no experimental information on the Dalitz-plot distribution of the decay $\eta' \rightarrow \pi^+\pi^-\pi^0$, nor a measurement of $\mathcal{B}(\eta' \rightarrow \pi^0\pi^0\pi^0)$. Therefore, we make use of the fact that the branching fraction $\mathcal{B}(\eta' \rightarrow \pi^+\pi^-\pi^0)$ depends on the coupling constant $g_{\eta'\rho\pi}$, under the conservative assumption that the $\rho^\pm$ states dominate the decay $\eta' \rightarrow \pi^+\pi^-\pi^0$. This will yield a conservative upper bound on $g_{\eta'\rho\pi}$, from which we obtain an upper bound on the ρ contribution to $\tau \rightarrow \pi\eta'\nu_\tau$. We discuss the likelihood of this assumption and its implications below.

The differential branching fraction of $\eta' \rightarrow \pi^+\pi^-\pi^0$ as a function of the Dalitz-plot position is given by

$$\frac{d\Gamma_{\eta' \rightarrow \pi^+\pi^-\pi^0}}{\Gamma_{\eta'}} = \frac{(g_{\eta'\rho\pi}g_{\rho\pi\pi})^2}{384\sqrt{3}\pi^3} \frac{Q^2}{m_{\eta'}\Gamma_{\eta'}} |\tilde{\mathcal{M}}|^2 dX dY, \quad (4)$$

where

$$Q \equiv m_{\eta'} - 3m_\pi \quad (5)$$

is the kinetic energy in the decay, and

$$\begin{aligned} X &\equiv \frac{\sqrt{3}}{Q}(T_+ - T_-), \\ Y &\equiv \frac{3}{Q}T_0 - 1 \end{aligned} \quad (6)$$

are the Dalitz-plot variables, with T_c being the kinetic energy of the pion with charge c . Assuming ρ dominance, we obtain from Eq. (15) of Ref. [1] the reduced matrix element

$$\tilde{\mathcal{M}} = -2 \frac{rY - \frac{1}{3}r^2(Y^2 + X^2)}{1 - \frac{2}{3}rY + \frac{1}{3}r^2(\frac{1}{3}Y^2 - X^2)}, \quad (7)$$

where

$$r = \frac{m_{\eta'}Q}{m_\rho^2 - \frac{1}{3}m_{\eta'}^2 - m_\pi^2 - i\Gamma_\rho m_\rho} = 1.6 + 0.7i. \quad (8)$$

The product $(g_{\eta'\rho\pi}g_{\rho\pi\pi})^2$ is then found by integrating Eq. (4) over the Dalitz plot. In the $\eta \rightarrow \pi^+\pi^-\pi^0$ case, we exploited the small value of r to simplify the expression by expanding in r . Due to the $O(1)$ value of r for $\eta' \rightarrow \pi^+\pi^-\pi^0$, we resort to numerical integration, which yields

$$\int |\tilde{\mathcal{M}}|^2 dX dY = 2.4. \quad (9)$$

From this we obtain, using $\mathcal{B}(\eta' \rightarrow \pi^+\pi^-\pi^0) = 3.7 \times 10^{-3}$ [9] and $g_{\rho\pi\pi} = 6.0$ [1],

$$g_{\eta'\rho\pi} < 0.025. \quad (10)$$

As a cross check, we apply the procedure to the decay $\eta \rightarrow \pi^+\pi^-\pi^0$, obtaining $g_{\eta\rho\pi} < 0.52$. This value is to be compared to the one obtained from the more precise Dalitz-plot analysis in Ref. [1], $g_{\eta\rho\pi} \approx 0.085$. The factor of 6 ratio between the results reflects the fact that the procedure used here yields but a conservative upper bound, obtained by assuming that the decay $\eta' \rightarrow \pi^+\pi^-\pi^0$ is dominated by the $\rho^\pm$ resonances. This assumption is manifestly false, as the $\eta' \rightarrow \pi^+\pi^-\pi^0$ Dalitz-plot distribution is in much better agreement with a flat distribution than with that expected from $\rho^\pm$ dominance [9]. By contrast, in Ref. [1], the value of $g_{\eta\rho\pi}$ obtained from the Dalitz-plot distribution yielded good agreement between the expected and measured values of $\mathcal{B}(\eta \rightarrow \pi^+\pi^-\pi^0)$.

With this point in mind, we proceed to use the upper bound on $g_{\eta'\rho\pi}$ to calculate the upper bound on the ρ contribution to $\mathcal{B}(\tau \rightarrow \pi\eta'\nu_\tau)$. We do this by relating $\mathcal{B}(\tau \rightarrow \rho(\pi\eta')\nu_\tau)$ to $\mathcal{B}(\tau \rightarrow \rho(\pi\eta)\nu_\tau)$ via the ratio of coupling constants and phase-space factors

$$\frac{\mathcal{B}(\tau \rightarrow \rho(\pi\eta')\nu_\tau)}{\mathcal{B}(\tau \rightarrow \rho(\pi\eta)\nu_\tau)} \approx \left(\frac{g_{\eta'\rho\pi}}{g_{\eta\rho\pi}} \right)^2 \frac{V(\tau \rightarrow \rho(\pi\eta')\nu_\tau)}{V(\tau \rightarrow \rho(\pi\eta)\nu_\tau)}, \quad (11)$$

where $\rho(\pi\eta')$ indicates that the ρ is observed in the $\pi\eta'$ final state, and $V(X)$ is the integral over the Dalitz plot of the three-body decay X . The ratio of phase-space integrals is 0.06, with up to 15% variation depending on whether one uses Blatt-Weisskopf and s -dependent widths for the ρ and on the choice of angular distribution. Using $\mathcal{B}(\tau \rightarrow \rho(\pi\eta)\nu_\tau) = 3.6 \times 10^{-6}$ [1], we obtain

$$\mathcal{B}(\tau \rightarrow \rho(\pi\eta')\nu_\tau) < 2 \times 10^{-8}, \quad (12)$$

more than two orders of magnitude below the *BABAR* upper limit, Eq. (3).

Next, we evaluate the contribution of the on-shell ρ' . One expects that this state, being a radial excitation and hence having a node in its wave-function, couples to the ground-state particles η and π more weakly than

the ρ . We hypothesize that this ρ' suppression mechanism works equally strongly for the final states $\pi\eta'$ and $\pi\pi$, leading to an equality of the ratios of the squared matrix elements

$$\begin{aligned} & \frac{\mathcal{B}(\tau \rightarrow \rho'_{(\pi\eta')}\nu_\tau)}{\mathcal{B}(\tau \rightarrow \rho_{(\pi\eta')}\nu_\tau)} \frac{V(\tau \rightarrow \rho_{(\pi\eta')}\nu_\tau)}{V(\tau \rightarrow \rho'_{(\pi\eta')}\nu_\tau)} \\ & \approx \frac{\mathcal{B}(\tau \rightarrow \rho'_{(\pi\pi)}\nu_\tau)}{\mathcal{B}(\tau \rightarrow \rho_{(\pi\pi)}\nu_\tau)} \frac{V(\tau \rightarrow \rho_{(\pi\pi)}\nu_\tau)}{V(\tau \rightarrow \rho'_{(\pi\pi)}\nu_\tau)}. \end{aligned} \quad (13)$$

The relevant phase-space integral ratios are

$$\begin{aligned} & \frac{V(\tau \rightarrow \rho_{(\pi\eta')}\nu_\tau)}{V(\tau \rightarrow \rho'_{(\pi\eta')}\nu_\tau)} \approx 0.06, \\ & \frac{V(\tau \rightarrow \rho_{(\pi\pi)}\nu_\tau)}{V(\tau \rightarrow \rho'_{(\pi\pi)}\nu_\tau)} \approx 2.5. \end{aligned} \quad (14)$$

We use the upper bound of Eq. (12) and the central value plus one standard deviation of the recent Belle result [10]

$$\sqrt{\frac{\mathcal{B}(\tau \rightarrow \rho'_{(\pi\pi)}\nu_\tau)}{\mathcal{B}(\tau \rightarrow \rho_{(\pi\pi)}\nu_\tau)}} = 0.15 \pm 0.05^{+0.15}_{-0.04} \quad (15)$$

to obtain the conservative upper limit

$$\mathcal{B}(\tau \rightarrow \rho'_{(\pi\eta')}\nu_\tau) < 8 \times 10^{-8}. \quad (16)$$

We note that this is an upper bound both due to the way we use Eq. (15) and since Eq. (12) is an upper bound.

III. THE $L = 0$ CONTRIBUTION

Calculating the $L = 0$ contributions to $\mathcal{B}'$ is not as straightforward as the $L = 1$ case, where one can make use of the dominant ρ coupling to the leptonic vector current. Therefore, it is important to evaluate the scalar component using different methods, as has been done for the $\tau \rightarrow \pi\eta\nu_\tau$ decay [1, 2, 3]. It should be noted that these calculations are performed under the assumption that the relevant scalar resonances are $\bar{u}d$ states. The coupling of a 4-quark state to the $\bar{u}d$ scalar current is ‘‘Zweig-Rule’’ suppressed, making it significantly smaller than the predictions.

Here we perform a more detailed version of the calculation used in Ref. [1]. We begin with the ratio of branching fractions

$$\begin{aligned} R_{a_1}^{a_0} & \equiv \frac{\mathcal{B}(\tau \rightarrow a_0\nu_\tau)}{\mathcal{B}(\tau \rightarrow a_1\nu_\tau)} \\ & = \frac{p_{a_0}}{p_{a_1}} \times \frac{|\langle a_0 | V_{h\mu} | 0 \rangle \langle \nu_\tau | J_l^\mu | \tau \rangle|^2}{|\langle a_1 | A_{h\mu} | 0 \rangle \langle \nu_\tau | J_l^\mu | \tau \rangle|^2}, \end{aligned} \quad (17)$$

where a_0 stands for either $a_0(980)$ or $a_0(1450)$, a_1 is the $a_1(1260)$, p_X is the τ -rest-frame momentum of the products of the decay $\tau \rightarrow X\nu_\tau$, $V_{h\mu} \equiv \bar{\psi}_u(x)\gamma_\mu\psi_d(x)$ is the hadronic vector current, $A_{h\mu} \equiv \bar{\psi}_u(x)\gamma_\mu\gamma^5\psi_d(x)$ is the

hadronic axial vector current, and $J_l^\mu \equiv \bar{\psi}_{\nu_\tau}(x)\gamma^\mu(1 - \gamma^5)\psi_\tau(x)$ is the leptonic current. The calculation of the leptonic parts of this ratio is well defined, while all the uncertainty in the hadronic parts comes down to a single parameter ξ , which shall be defined shortly. With this in mind, we can take the a_0 matrix element to be

$$\langle a_0 | V_{h\mu} | 0 \rangle = f_0 \frac{q_\mu}{m_{a_0}} \langle a_0 | S_h | 0 \rangle, \quad (18)$$

where f_0 is an isospin-violation suppression factor, and $S_h \equiv \bar{\psi}_u(x)\psi_d(x)$ is the scalar current operator. The weak vector current is conserved up to the difference between the u - and d -quark masses, plus a smaller electromagnetic part that we neglect. Therefore,

$$\partial^\mu V_{h\mu} \approx (m_d - m_u)S_h. \quad (19)$$

Using this relation in Eq. (18) yields

$$f_0 = \frac{m_d - m_u}{m_{a_0}}. \quad (20)$$

We use the fact that both the a_0 and the a_1 are P -wave states to relate the axial and scalar decay constants

$$\langle a_1 | A_\mu | 0 \rangle = \xi \epsilon_\mu^* \langle a_0 | S | 0 \rangle. \quad (21)$$

We note that this is reminiscent of applying $SU(6)$ [11] or, in this case, just $SU(4)$ [12] flavor-spin symmetry to the ($L = 0$) 15-plet plus singlet containing the π , ρ , η , and ω , or the ($L = 1$) states a_0 , a_1 , f_0 , and h_1 .

Naively, one expects ξ in Eq. (21) to be of order unity. However, this parameter incorporates all the hadronic uncertainty in our procedure. With Eqs. (18-21), Eq. (17) becomes, after spin averaging and index contraction,

$$\begin{aligned} R_{a_1}^{a_0} & = |\xi|^2 \frac{p_{a_0}}{p_{a_1}} \left(\frac{m_d - m_u}{m_{a_0}} \right)^2 \\ & \times \frac{m_\tau^2 - m_{a_0}^2}{m_\tau^2 - m_{a_1}^2} \left(\frac{m_{a_1}}{m_{a_0}} \right)^2 \frac{1}{1 + 2(m_\tau/m_{a_1})^2}. \end{aligned} \quad (22)$$

This yields the branching fractions

$$\begin{aligned} \mathcal{B}(\tau \rightarrow a_0(980)\nu_\tau) & = 1.6 \times 10^{-6} |\xi|^2, \\ \mathcal{B}(\tau \rightarrow a_0(1450)\nu_\tau) & = 6.4 \times 10^{-8} |\xi|^2, \end{aligned} \quad (23)$$

where, as in Ref. [1], we chose the mass difference of the two light quarks to be 4 MeV [13] and, assuming that the $\tau \rightarrow 3\pi\nu_\tau$ decay is dominated by the a_1 , we took $\mathcal{B}(\tau \rightarrow a_1\nu_\tau) = 0.18$. We compare $\mathcal{B}(\tau \rightarrow a_0(980)\nu_\tau)$ of Eq. (23) with the value $\mathcal{B} = 1.2 \times 10^{-5}$, obtained from the more elaborate calculation of Ref. [2], minus the ρ contribution to $\mathcal{B}$, which is 3.6×10^{-6} [1]. This yields $|\xi|^2 \approx 5$, from which we conclude

$$\mathcal{B}(\tau \rightarrow a_0(1450)\nu_\tau) \approx 3 \times 10^{-7}. \quad (24)$$

The $a_0(1450)$ contribution to $\tau \rightarrow \pi\eta'\nu_\tau$ depends also on the branching fraction $\mathcal{B}(a_0(1450) \rightarrow \pi\eta')$, regarding

which there is only partial information. However, from the branching-fraction measurements that have been made [13], it is clear that $\mathcal{B}(a_0(1450) \rightarrow \pi\eta') < 0.3$. Hence

$$\mathcal{B}(\tau \rightarrow a_0(1450)_{(\pi\eta')} \nu_\tau) < 1 \times 10^{-7}. \quad (25)$$

If the $a_0(1450)$ is a radial excitation, which is the case if the $a_0(980)$ is the $\bar{u}d$ ground state, then $\mathcal{B}(\tau \rightarrow a_0(1450)_{(\pi\eta')} \nu_\tau)$ should be suppressed by an additional wave-function overlap factor.

Next, we look at the contribution of the $a_0(980)$ to $\tau \rightarrow \pi\eta' \nu_\tau$, which can be extracted from the relation

$$\frac{\mathcal{B}(\tau \rightarrow \nu a_0(980)_{(\pi\eta')})}{\mathcal{B}(\tau \rightarrow \nu a_0(980)_{(\pi\eta)})} = \frac{V(\tau \rightarrow \nu a_0(980)_{(\pi\eta')})}{V(\tau \rightarrow \nu a_0(980)_{(\pi\eta)})} R_{\eta'}^{\eta'}, \quad (26)$$

where

$$R_{\eta'}^{\eta'} \equiv \left| \frac{\mathcal{M}(a_0(980) \rightarrow \pi\eta')}{\mathcal{M}(a_0(980) \rightarrow \pi\eta)} \right|^2 \quad (27)$$

is the square of the ratio between the relevant hadronic-decay matrix elements. We assume that $R_{\eta'}^{\eta'}$ equals the corresponding ratio of $a_0(1450)$ -decay matrix elements, and is hence obtained from

$$R_{\eta'}^{\eta'} \approx \frac{\mathcal{B}(a_0(1450) \rightarrow \pi\eta')}{\mathcal{B}(a_0(1450) \rightarrow \pi\eta)} \times \frac{p_\eta}{p_{\eta'}}, \quad (28)$$

where p_X is the $a_0(1450)$ -rest-frame momentum of the products of the decay $a_0(1450) \rightarrow \pi X$. Given the $\sim 50\%$ error [13] on the ratio of branching fractions appearing in Eq. (28) and the uncertainty on the $a_0(1450)$ width, $R_{\eta'}^{\eta'}$ comes out in the range $[0.25, 1.25]$. The ratio of the phase-space integrals in Eq. (26) is 0.06, with some dependence on what one takes for the $a_0(980)$ width. Using

the range for $\mathcal{B}$ from Eq. (2), we obtain

$$\mathcal{B}(\tau \rightarrow a_0(980)_{(\pi\eta')} \nu_\tau) \approx [0.2 \text{ to } 1.2] \times 10^{-6}. \quad (29)$$

IV. CONCLUSIONS

Combining Eqs. (12), (16), (25), and (29), we obtain the branching fraction limit

$$\mathcal{B}(\tau \rightarrow \pi\eta' \nu_\tau) < 1.4 \times 10^{-6}, \quad (30)$$

in no conflict with the experimental upper limit, Eq. (3), which is about five times greater. Our result is dominated by the $a_0(980)$ contribution, assuming it is a $\bar{u}d$ state.

The experimental limit was obtained with only a third of the currently available *BABAR* and *Belle* data sets, and with the η reconstructed only in the $\gamma\gamma$ final state. Therefore, an improvement in the limit can be expected from the current generation of *B* factories, but probably not to the level of Eq. (30). By contrast, a Super *B* factory [14], with two orders of magnitude more luminosity, will be able to use $\mathcal{B}$ and $\mathcal{B}'$ to investigate the nature of the $a_0(980)$ and to search for new interactions mediated by heavy scalars [1].

Acknowledgments

This research was supported in part by grant number 2006219 from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel. The authors thank Leonid Frankfurt and Swagato Banerjee for useful suggestions and discussions.

-
- [1] S. Nussinov and A. Soffer, Phys. Rev. D **78**, 033006 (2008) [arXiv:0806.3922 [hep-ph]].
 - [2] H. Neufeld and H. Rupertsberger, Z. Phys. C **68**, 91 (1995).
 - [3] A. Pich, Phys. Lett. B **196**, 561 (1987); S. Tisserant and T. N. Truong, Phys. Lett. B **115**, 264 (1982);
 - [4] J. E. Bartelt *et al.* [CLEO Collaboration], Phys. Rev. Lett. **76**, 4119 (1996).
 - [5] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **77**, 112002 (2008) [arXiv:0803.0772 [hep-ex]].
 - [6] R. Escribano, arXiv:0807.4201 [hep-ph]; F. Ambrosino *et al.* [KLOE Collaboration], Phys. Lett. B **648**, 267 (2007) [arXiv:hep-ex/0612029].
 - [7] R. L. Jaffe, Phys. Rev. D **15**, 267 (1977).
 - [8] G. 't. Hooft, G. Isidori, L. Maiani, A. D. Polosa and V. Riquer, arXiv:0801.2288 [hep-ph].
 - [9] P. Naik *et al.* [CLEO Collaboration], Phys. Rev. Lett. **102**, 061801 (2009) [arXiv:0809.2587 [hep-ex]].
 - [10] M. Fujikawa *et al.* [Belle Collaboration], Phys. Rev. D **78**, 072006 (2008) [arXiv:0805.3773 [hep-ex]].
 - [11] F. Gursey and L. A. Radicati, Phys. Rev. Lett. **13**, 173 (1964).
 - [12] E. Wigner, Phys. Rev. **51**, 106 (1937).
 - [13] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
 - [14] M. Bona *et al.*, arXiv:0709.0451 [hep-ex].